

POISON
 KEEP OUT OF REACH OF CHILDREN
 READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Haloxyfop 520EC®

Herbicide

ACTIVE CONSTITUENT: 520 g/L HALOXYFOP present as the HALOXYFOP-P METHYL ESTER

GROUP A HERBICIDE

For the post-emergent control of a wide range of annual and perennial grass weeds in grain legume and oilseed crops, lucerne, medic and clover pasture and seed crops, forestry, bananas, citrus, grapes, pineapples, pome and stone fruit, pyrethrum, tropical fruit and nut crops as specified in the Directions for Use.

IMPORTANT: READ THIS LEAFLET BEFORE OPENING OR USING THIS PRODUCT

Sabakem Pty Ltd ABN 34 151 682 138
 Suite 809, Level 8, 2 Queen Street, Melbourne VIC 3000
 Tel: 03 9629 3979 • Fax: 1300 242 436
www.sabakem.com

Leaflet_1117

APVMA Approval No: 69155/60090

DIRECTIONS FOR USE

RESTRAINTS:

DO NOT apply to weeds which may be stressed (not actively growing) due to prolonged periods of extreme cold, moisture stress (water-logged or drought affected), poor nutrition or previous herbicide treatment as reduced levels of control may result.

DO NOT spray if rain is likely to occur within one hour.

Table 1a: Winter Crops – Canola, Chickpeas, Faba beans, Field peas, Lentils, Linola, Linseed Lupins, Lucerne, Vetch, Medic and Clover pastures or seed crops				
WEEDS CONTROLLED	WEED GROWTH STAGE	RATE		CRITICAL COMMENTS
		With Uptake ¹ Spraying Oil	With a Non-ionic Wetter ²	
Annual Ryegrass	2 to 4 leaf	75 mL/ha	100 mL/ha	<p>CANOLA, LINOLA AND LINSEED DO NOT apply after the 8 leaf stage of the crop. DO NOT apply after the commencement of stem elongation. This means that application must not occur after the 8 leaf stage, or if stem elongation commences before the 8 leaf stage, application must not occur after stem elongation has commenced. DO NOT apply more than 1 application of herbicide containing haloxyfop per crop. DO NOT apply after grazing.</p> <p>^{1 2} See GENERAL INSTRUCTIONS Spraying oils/wetters section.</p> <p>FIELD PEAS AND CANOLA The only oil recommended for use with Sabakem Haloxyfop 520EC® Herbicide is Uptake¹ Spraying Oil. Sabakem Haloxyfop 520EC® Herbicide + Lontrel 750SG + Uptake¹ Spraying Oil are compatible and selective to canola. This tank mix is also compatible with atrazine and selective to triazine tolerant canola.</p> <p>LUPINS AND FIELD PEAS Mixtures with Brodal[†] or simazine may cause crop yellowing and separate applications are recommended.</p> <p>CHICKPEAS, FABA BEANS, LENTILS AND VETCH, LINOLA, LINSEED Broadleaf herbicides should not be added to Sabakem Haloxyfop 520EC® Herbicide. Apply Sabakem Haloxyfop 520EC® Herbicide and broadleaf herbicides at least a week apart.</p> <p>LUCERNE, CLOVER OR MEDIC PASTURES: If grazed or cut for hay immediately prior to treatment, delay application until all grasses have fully expanded leaves. Use 75 mL + spraying oil or 100 mL + wetter/ha. (See GENERAL INSTRUCTIONS, Spraying Oils/wetters section). If silver grass (<i>Vulpia</i> spp.) is present in pasture, simazine should be tank mixed with the higher rate of Sabakem Haloxyfop 520EC® Herbicide plus a non-ionic wetter.</p>
	Early tillering	100 mL/ha	100 mL/ha	
Barley Grass	2 to 4 leaf	50 mL/ha	75 mL/ha	
	Early tillering	75 mL/ha	100 mL/ha	
Brome Grass		Early tillering	75 mL/ha	
	Paradoxa Grass			
Volunteer cereals	2 to 4 leaf	37.5 mL/ha	50 mL/ha	
WA, SA, Vic, Tas, Southern and Central NSW	Early tillering	50 mL/ha	75 mL/ha	
Northern NSW & Qld	Early tillering	75 mL/ha	100 mL/ha	
				Northern NSW & Qld

Table 1b: Winter crop growth stage application windows	
Crop	Crop Growth Stage
Lucerne, Medic and Clover pastures or Seed crops	Apply from 2nd trifoliolate leaf onwards. For <i>Erodium</i> spp., spraying, apply from cotyledon crop stage onwards.
Canola, Linola and Linseed	Apply from 2 leaf to 8 leaf stage of crop growth. DO NOT apply after the commencement of stem elongation. This means that application must not occur after the 8 leaf stage, or if stem elongation commences before the 8 leaf stage, application must not occur after stem elongation has commenced.
Chickpeas, Faba beans, Field peas, Lentils, Lupins and Vetch	Apply from 2nd leaf, 2nd node or 2nd branch prior to flowering.

Table 2a: Lucerne, Medic and Clover seed crop s and pastures. See table 1b for crop stages.			
WEEDS CONTROLLED	WEED GROWTH STAGE	RATE With Uptake ¹ Spraying Oil	CRITICAL COMMENTS
Prairie Grass (<i>Bromus catharticus</i>)	Up to early tillering	100 mL/ha	<p>¹ See GENERAL INSTRUCTIONS, Spraying oils/wetters section.</p> <p>³ Use lower rate when growing conditions and crop or pasture competition are good and when weed populations are below 100 plants/m². Use the higher rate when weed populations are above 100 plants/m² or when crop or pasture competition is poor.</p> <p>Note: Storksbill may not be controlled if simazine or Broadstrike[†] are tank-mixed with Sabakem Haloxyfop 520EC® Herbicide.</p> <p>LUCERNE, CLOVER OR MEDIC PASTURES If grazed or cut for hay immediately prior to treatment, delay application until all grasses have fully expanded leaves. Use 75 mL + spraying oil or 100 mL + wetter/ha. (See GENERAL INSTRUCTIONS, Spraying Oils/Wetter section). If Silver Grass (<i>Vulpia</i> spp.) is present in pasture, simazine should be tank-mixed with the higher rate of Sabakem Haloxyfop 520EC® Herbicide plus a non-ionic wetter.</p> <p>⁴ For best suppression of couch or control of Rhodes Grass, make two (2) applications of Sabakem Haloxyfop 520EC® Herbicide 2 to 4 weeks apart. Time second application to coincide with tillering stage of weeds and just after irrigation or significant rain. Only treat actively growing weeds which are not moisture stressed. Use these rates for control of couch and Rhodes grass.</p>
Musky or Ferny Leaf Storksbill (<i>Erodium moschatum</i>)	Up to 6 leaf or 5 cm diameter	50 to 75 mL/ha ³	
Common Crowsfoot or Common Storksbill (<i>Erodium cicutarium</i>)	Up to 6 leaf or 5 cm diameter	50 to 75 mL/ha ³	
Long or Shiny Leaf Storksbill (<i>E. botrys</i>)	Up to 8 leaf or 5 cm diameter	75 to 100 mL/ha	
Couch Grass (suppression)	Tillering seedlings	150 mL/ha + 150 mL/ha ⁴	
Rhodes Grass (control)			
Couch Grass (control)	Established stands	400 to 800 mL/ha	
Rhodes Grass (control)			

Table 3a: Summer crops – Cotton, Cowpea, Lucerne, Mung beans, Navy beans, Peanuts, Soybeans, Sunflowers			
WEEDS CONTROLLED	WEED GROWTH STAGE	RATE With Uptake ¹ Spraying Oil	CRITICAL COMMENTS
Australian Millet	2 leaf to tillering up to 15 cm	150 mL/ha	<p>¹ See GENERAL INSTRUCTIONS, Spraying Oils/wetters section.</p> <p>NAVY BEANS, PEANUTS, SOYBEANS: For broadleaf weed control, Sabakem Haloxyfop 520EC® Herbicide at 150 mL/ha plus wetter may be tank mixed with Blazer[†] (except on Navy beans) or Basagran[†]. Tank mixtures may cause transient leaf spotting on the crop but do not normally affect yield. DO NOT tank mix broadleaf herbicides with Sabakem Haloxyfop 520EC® Herbicide if grasses have begun tillering or if the grasses are under moisture stress. DO NOT add Uptake¹ Spraying Oil when mixing with Blazer[†] or Basagran[†]. DO NOT use Blazer[†] or Basagran[†] tank-mixes on cowpea.</p>
Barnyard Grass	2 to 5 leaf	100 mL/ha	
	Tillering up to 15 cm	150 mL/ha	
Crowsfoot Grass	2 leaf to tillering up to 15 cm	150 mL/ha	
		Green Panic	
Johnson Grass (rhizome)	2 to 5 leaf	100 mL/ha	
Johnson Grass (seedling)	Tillering and up to 15 cm	150 mL/ha	
Liverseed Grass (seedling)			
Mossman River Grass	2 to 5 leaf	100 mL/ha	
Summer Grass	2 leaf to tillering up to 15 cm	150 mL/ha	
		Volunteer Cereals	
Volunteer Cereals	2 to 4 leaf	100 mL/ha	
	Tillering up to 15 cm	150 mL/ha	

Table 3b: Summer crop growth stage application windows	
Crop	Crop Growth Stage
Lucerne	Apply from 2nd trifoliolate leaf onwards.
Cowpea, Mung beans, Navy beans, Soybeans	Apply from 2nd leaf to flowering.
Peanuts	Apply from 2nd leaf to pegging.
Cotton	Apply from 2nd leaf to before the onset of flowering.
Sunflowers	Apply from 2nd leaf to head initiation.

CROPS	CROP GROWTH STAGE	WEEDS CONTROLLED	WEED GROWTH STAGE	RATE With Uptake [†] Spraying Oil [†]	CRITICAL COMMENTS
Orchard, Vine and Plantation Crops, including: Apples, Avocado, Banana, Blueberry, Citrus, Custard apple, Feijoa, Grapevines, Guava, Kiwifruit, Litchi (Lychee), Longan, Mango, Nashi, Nut trees, Passionfruit, Paw paw, Pear, Persimmon, Pineapple, Rambutan, Stone fruit	All growth stages	Perennial grasses: Couch Rhodes Grass Slender Rats Tail Grass	Established stands	400 to 800 mL/ha	¹ See GENERAL INSTRUCTIONS, Spraying Oils/Wetter section. Spray should be directed to the base of the tree or vine avoiding contact with fruit and foliage. Spot spray: Use 25 mL to 50 mL/100 L of water. Use higher rate on late tillering mature grasses. Annual Grasses: Where treated in association with perennial grasses, these annual grasses will be controlled.
		Buffel Grass Green panic Johnson grass Kikuyu <i>Paspalum</i> spp. <i>Setaria</i> spp.	Vegetative to early tillering	200 mL/ha	
		Annual grasses: Annual Ryegrass Barley Grass Barnyard Grass Brome Grass Crowsfoot Grass Lesser Canary Grass Liverseed grass Mossman River Grass Paradoxa Grass Summer Grass Volunteer Cereals Wild Oats	Late tillering	400 mL/ha	
Forestry: <i>Pinus radiata</i> , <i>Eucalyptus</i> spp.	All growth stages	Annual grasses: As above	2 leaf to tillering	200 mL/ha	Annual Grasses: Where treated in association with perennial grasses, these annual grasses will be controlled.
		Annual grasses: As above	Vegetative to tillering	125 to 250 mL/ha	
Forestry: <i>Pinus pineaster</i>	All growth stages	Annual grasses: As above	Vegetative to tillering	125 to 250 mL/ha	Forestry: For annual grasses apply lowest rate to newly emerged grasses, increasing the rate as they develop.
Pyrethrum	All growth stages	Barley Grass Brome Grass Rope Twitch Barnyard Grass <i>Erodium</i> spp. Volunteer cereals	Vegetative to tillering	100 to 250 mL/ha	Pyrethrum Tasmania only: For <i>Erodium</i> spp., apply 75 to 100 mL/ha if the main weed is <i>E. botrys</i> . Use 50 to 75 mL/ha if either <i>E. cicutarium</i> or <i>E. moschatum</i> are the main weeds.

GENERAL INSTRUCTIONS

MIXING

- Add water to the spray tank to 10 cm above the level of agitation and ensure the agitation device is working vigorously. (There must be a minimum of 100 L of water in the tank before any pesticide is added.)
- If tank mixing, firstly add any soluble liquid formulations (eg: Lontrel[†] Herbicide) and allow agitation for approximately one minute.
- Then add Sabakem Haloxyfop 520EC[®] Herbicide at the point where agitation is strongest. (DO NOT add Sabakem Haloxyfop 520EC[®] Herbicide through a strainer or sieve). Allow further agitation for one minute.
- Half fill the spray tank.
- If using wettable powder or water dispersible granules, or other emulsifiable concentration formulations (eg: Lorsban[†] 750WG or Le-Mat[†], these should be added after the Sabakem Haloxyfop 520EC[®] Herbicide to the half-full spray tank ensuring vigorous agitation.
- Finally add Uptake[†] Spraying Oil or approved alternate spraying oil/wetter. (See section on spraying oils/wetters) and continue filling the tank to the required volume maintaining agitation at all times.
- Only mix sufficient solution for immediate use. Sabakem Haloxyfop 520EC[®] Herbicide and any other tank mixes should be applied immediately for best results.

Spraying Oils/wetters

- Spraying Oils:** It is essential to add an adjuvant to Sabakem Haloxyfop 520EC[®] Herbicide. Best results will be achieved with Uptake[†] Spraying Oil at 0.5 L/100 L of spray solution. Alternatively, other oils plus a non-ionic wetter may also be used. When other crop spraying oils are used, mix at 1.0 L/100 L and add a non-ionic wetter (surfactant) at 200 mL/100 L of spray solution. Use of an oil is not always recommended. See Critical Comments for specific situation recommendations.
- Non-ionic Wetters:** When Uptake[†] or other oils are not used, a 100% concentrate non-ionic wetting agent such as BS-1000[†] at 200 mL/100 L must be used along with the higher rate of Sabakem Haloxyfop 520EC[®] Herbicide as specified in the Directions for Use.

Where water volumes of less than 50 L/ha are used, DO NOT use less than 250 mL/ha of Uptake[†] or 500 mL/ha for oils other than Uptake[†] or less than 100 mL/ha of wetter.

CANOLA, LUCERNE, MEDIC AND CLOVER PASTURES AND SEED CROPS

When tank mixing Sabakem Haloxyfop 520EC[®] Herbicide with Lontrel[†] herbicides (canola only) or Broadstrike[†] (lucerne, clover and medics), use Uptake[†] Spraying Oil with the lower rates of Sabakem Haloxyfop 520EC[®] Herbicide or a wetting agent with the higher rates of Sabakem Haloxyfop 520EC[®] Herbicide unless otherwise specified. When mixing Sabakem Haloxyfop 520EC[®] Herbicide with other broadleaf herbicides on these crops, DO NOT use an oil, use a wetter instead.

FIELD PEAS AND CANOLA

The oil recommended is Uptake[†] Spraying Oil. Hasten[†] is also recommended for use with tank mixtures of Sabakem Haloxyfop 520EC[®] Herbicide and Select[†] Herbicide.

For canola, Sabakem Haloxyfop 520EC[®] Herbicide + Lontrel[†] 750SG + Uptake[†] Spraying Oil are compatible and selective to canola. This tank mixture is also compatible with atrazine or simazine and selective to triazine tolerant canola.

NAVY BEANS, PEANUTS, SOYBEANS

When mixing with Blazer[†] or Basagran[†], DO NOT add spraying oil to these mixtures. DO NOT use these tank mixes on cowpea.

COMPATIBILITY

Ground use only: Sabakem Haloxyfop 520EC[®] Herbicide can be tank mixed with:

Insecticides: Dimethoate, Endosulfan, Chlorpyrifos 500 EC Insecticide, Lorsban[†] 750 WG Insecticide, Omethoate

Herbicides: Atrazine, Basagran[†], Blazer[†], Broadstrike[†] Herbicide, Clopyralid Herbicide, Lontrel[†] 750SG, MCPA ester (LVE) – DO NOT exceed 700 mL/ha of MCPA LVE, Oryzalin, Clethodim Herbicide, Simazine, Fluroxypyr 200 Herbicide

Fungicides: Dithane DF[†], Dithane Rainshield

Trace Elements: Magnesium sulphate, Zinc sulphate

Sabakem Haloxyfop 520EC[®] Herbicide is NOT COMPATIBLE with 2,4-D or MCPA as sodium or amine salts.

Aerial use: No product, other than a recommended crop oil or wetter, should be mixed with Sabakem Haloxyfop 520EC[®] Herbicide when applied by air, except for addition of Lontrel[†] Forestry Herbicide for use in forestry and Lontrel 750SG for use in canola only.

APPLICATION

Apply Sabakem Haloxyfop 520EC[®] Herbicide in sufficient water to obtain good coverage. It should be applied by an accurately calibrated ground rig or aircraft delivering droplets with a VMD of 200 to 300 microns.

The following spray volumes are recommended:

Ground application:	50 to 150 L/ha
Aerial application:	30 L/ha minimum

Use higher water volumes in orchards and in dense crops where the weeds may be shielded by the crop canopy.

CLEANING SPRAY EQUIPMENT

If broadleaf herbicides, particularly sulfonylureas, have been used in the spray equipment at any time prior to Sabakem Haloxyfop 520EC[®] Herbicide, particular care should be taken to follow the directions on the relevant broadleaf herbicide label for equipment cleaning, or damage to susceptible crops may occur.

After using Sabakem Haloxyfop 520EC[®] Herbicide, empty the tank completely and drain the whole system. Thoroughly wash inside the tank using a pressure hose, drain the tank and clean any filters in the tank, pump, line and nozzles.

To rinse: After cleaning the tank as above, quarter fill the tank with clean water and circulate through the pump, lines, hoses and nozzles. Drain and repeat the rinsing procedure twice.

To decontaminate: Before spraying cereals, maize, sorghum or other sensitive crops, wash the tank and rinse the system as above. Then, quarter fill the tank and add an alkali detergent (eg: Surf[®], Cold Water SURF Concentrate, DynamoMatic Concentrate, OMO[®] or DRIVE[®]) at 500 mL/100 L of water or the powder equivalent at 500 g/100 L of water, and circulate throughout the system for at least fifteen minutes. Drain the whole system. Remove filters and nozzles and clean them separately. Finally flush the system with clean water and allow to drain. Chlorine-based cleaners are not recommended.

Rinse water should be discharged onto a designated disposal area, or if this is unavailable, onto unused land away from desirable plants and water sources.

RESISTANT WEEDS WARNING

Sabakem Haloxyfop 520EC[®] Herbicide is a member of the aryloxyphenoxy propionate group of herbicides. Sabakem Haloxyfop 520EC[®] Herbicide has the acetyl CoA carboxylase inhibitor mode of action. For weed resistance management, Sabakem Haloxyfop 520EC[®] Herbicide is a Group A Herbicide. Some naturally-occurring weed biotypes resistant to Sabakem Haloxyfop 520EC[®] Herbicide and other inhibitors of acetyl CoA carboxylase herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Sabakem Haloxyfop 520EC[®] Herbicide or other inhibitors of acetyl CoA carboxylase. Since the occurrence of resistant weeds is difficult to detect prior to use, Sabakem Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant weeds.

GROUP A HERBICIDE

Strategies to minimize the risk of herbicide resistance are available. Contact your farm chemical supplier, consultant, local Department of Agriculture, or Sabakem representative.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Sabakem Haloxyfop 520EC[®] Herbicide damages cereals and grasses. DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. Cereal crops or grasses planted within twelve (12) weeks of application may be damaged by the residual effects of Sabakem Haloxyfop 520EC[®] Herbicide, particularly on light and red soils.

PROTECTION OF LIVESTOCK

DO NOT graze or cut treated crops for stock food except as specified under withholding periods.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Sabakem Haloxyfop 520EC[®] Herbicide is toxic to fish. DO NOT contaminate streams, rivers or waterways with the chemical or used container.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT store near feedstuffs, fertilisers or seeds. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SMALL SPILL MANAGEMENT

Wear protective equipment (See SAFETY DIRECTIONS). Apply absorbent material such as earth, sand, clay granules or cat litter to the spill. When absorption is complete, sweep up material and contain in a refuse vessel for disposal (see STORAGE AND DISPOSAL section). If necessary, wash the spill area with an alkali detergent and water and absorb the wash liquid for disposal as described above.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container, preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves, and face shield or goggles. After each day's use, wash gloves, face shield or goggles and contaminated clothing. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

SAFETY DATA SHEET

Additional information is listed on the Safety Data Sheet which is available from the supplier.

CONDITIONS OF SALE: The use of this product is beyond the control of Sabakem Pty Ltd. No warranty is expressed or implied regarding the suitability or efficiency for any purpose for which it is used by the buyer. Sabakem Pty Ltd accepts no responsibility for any consequences resulting from the use of this product. Sabakem Pty Ltd will not be held liable for any loss, injury or damage arising from the sale, supply or use of this product, whether through negligence or otherwise. No responsibility will be accepted for any consequences whatsoever resulting from the use of this product.

Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: Harmful if inhaled. Very toxic to aquatic life with long lasting effects.

© Registered trademarks of Sabakem Pty Ltd

† Not a Sabakem trademark

Table 5: Sabakem Haloxyfop 520EC[®] Herbicide and Clethodim Herbicide tank-mixes – Canola, Chickpeas, Faba beans, Field peas, Lupins, Lentils.

WEEDS CONTROLLED	WEED GROWTH STAGE	RATE		CRITICAL COMMENTS
		Sabakem Haloxyfop 520EC [®]	Clethodim Herbicide	
FOP/DIM susceptible Annual Ryegrass + Volunteer Barley Volunteer Wheat Brome Grass Wild Oats Barley Grass Phalaris	2 to 4 leaf	25 mL/ha	150 mL/ha	See GENERAL INSTRUCTIONS, Spraying Oils/wetters section. Use Uptake [†] Spraying Oil at 500 mL/100 L or Hasten [†] at 1 L/100 L. Apply at the same crop growth stages as those in Table 1b Winter Crops. Lentils: Apply up to 7 node-early branching crop growth stage only. Lupins: Not for Qld.
	Early tillering	38 mL/ha	150 mL/ha	
FOP resistant Annual ryegrass + Volunteer Barley Volunteer Wheat Brome Grass Wild Oats Barley Grass Phalaris	2 to 4 leaf	25 mL/ha	200 mL/ha	
	Early tillering	38 mL/ha	250 mL/ha	

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

HARVESTING WITHHOLDING PERIODS

NOT REQUIRED WHEN USED AS DIRECTED FOR:

Canola, Chickpeas, Cotton, Cowpea, Faba beans, Field Peas, Lentils, Linola, Linseed, Lupins, Mung beans, Navy beans, Orchard Crops, Peanuts, Plantation Crops, Soybeans, Sunflowers, Vetch or Vine Crops

DO NOT HARVEST FOR:

Medic and Clover seed crops: 7 DAYS AFTER APPLICATION

STOCK FOOD WITHHOLDING PERIODS

DO NOT GRAZE OR CUT FOR STOCK FOOD FOR:

Canola, Chickpeas, Cotton, Cowpea, Faba beans, Field Peas, Lentils, Linola, Linseed, Lupins, Mung beans, Navy beans, Peanuts, Soybeans, Sunflowers, Vetch: 28 DAYS AFTER APPLICATION

Lucerne: 21 DAYS AFTER APPLICATION

Medic and Clover Pasture: 7 DAYS AFTER APPLICATION

COTTON GIN TRASH MUST NOT BE FED TO ANIMALS.