



CROP	WEEDS CONTROLLED	WEED STAGE	RATE mL/ha	STATE	CRITICAL COMMENTS
<b>Wheat, barley, triticale, cereal rye</b> (including undersown with clover and/or Lucerne), and these cover crops in vineyards.  <b>Pasture</b> Clover and/or Lucerne based pasture (newly sown or established) including cover crops in vineyards	Wild radish	Up to 2 leaf stage and not more than 60 mm in diameter and where weed density is less than 50 plants/m <sup>2</sup>	350 mL	WA only	<b>CROP STAGE</b> <b>Cereals</b> 2 leaf to fully tillered (zadok's Z12-29). Optimum results are achieved when sprayed at 4-8 weeks post sowing.  <b>Warning:</b> Sabakem Reckon-B 275EC® Herbicide may cause transient crop yellowing of cereals. Refer to "Crop Tolerance section of General Instructions).  <b>Clover and Lucerne</b> Application is recommended prior to the 8th trifoliolate leaf stage. Application can be made from the 1st trifoliolate leaf stage in Qld, NSW, ACT and Vic only. In other States applications prior to the 3 leaf stage may result in crop damage if seedlings are under stress and in sandy soils. DO NOT apply to annual medics.  <b>Warning:</b> Sabakem Reckon-B 275EC® Herbicide may affect growth and seed set of some varieties of clover and Lucerne. (Refer to "Crop Tolerance" section of General Instructions).  <b>COVER CROPS IN VINEYARDS</b> When using in vineyard situations, apply during vine dormancy only. Contact with vines must be avoided. Particular care should be taken if applied in late autumn or early spring, when vines may not be fully dormant.  <b>WEED STAGE:</b> Apply from early post-emergence.  <b>APPLICATION:</b> Apply when weeds are actively growing. Ensure thorough coverage of weeds. Where crop or weed density is high, increase water volume.  In most situations the rate specified for each weed size will give satisfactory control. However, under certain conditions such as: * high crop and weed density; * late season germinations; * abnormal weed growth (including early flowering): higher rates of product (up to the maximum rate of application specified for that weed) may be required.  Sabakem Reckon-B 275EC® Herbicide will not effectively control: * regrowth of suppressed weeds; * transplanted weeds; * regrowth from rhizomes or roots; * weeds growing under stress from previous herbicide applications; * radish plants beyond rosette stage.  <b>WILD RADISH</b> Effective residual activity of this product may be reduced where: *rates lower than 1.0 L/ha are used: * dry conditions prevail; * poor coverage of the soil surface is achieved; * crop is grown in non-wetting sand; * soils have a high content of clay or organic matter.  <b>VOLUNTER LUPINS</b> In some situations, the higher rate of 1.0 L/ha may be required to effectively suppress volunteer lupins at the 4 leaf stage.  # Sabakem Reckon-B 275EC® Herbicide will suppress seedling dock but will not suppress regrowth from transplanted roots.
	Wild mustard, Wild radish	Up to 4 leaf stage and not more than 120 mm in diameter.	500 mL	All States	
	Canola (rapeseed) charlock, turnip weed, wild turnip	Up to 2 leaf stage and not more than 60 mm in diameter	500 mL	All States	
		Up to 4 leaf stage and not more than 120 mm in diameter.	750 mL		
	Shepherd's purse	Up to 6 leaf stage and not more than 150 mm in diameter.	750 ml	All States	
		Up to 8 leaf stage and not more than 180 mm in diameter.	1.0 L		
	Capeweed	Up to 2 leaf stage and not more than 60 mm in diameter	500 mL	All States	
		Up to 4 leaf stage and not more than 120 mm in diameter.	750 mL		
		Up to 6 leaf stage and not more than 150 mm in diameter.	1.0 L		
	Corn gromwell	Up to 2 leaf stage and not more than 60 mm in diameter	500 mL	All States	
		Up to 4 leaf stage and not more than 120 mm in diameter.	750 mL		
		Up to 6 leaf stage and not more than 150 mm in diameter.	1.0 L		
	Climbing buckwheat	Up to 2 leaf stage	500 mL	All States	
		Up to 4 leaf stage	750 mL		
		Up to 6 leaf stage	1.0 L		
	Deadnettle, Paterson's curse, (Salvation Jane), rough poppy	Up to 2 leaf stage	500 mL	All States	
		Up to 4 leaf stage	750 mL		
	Amsinckia				
	Doublegee (spiny emex)	Up to 2 leaf stage	500 mL	Qld, NSW, ACT, Vic, Tas, WA only	
		Up to 4 leaf stage	750 mL	All States	
Chamomile, common peppergrass, lesser swinecress, purple calandrinia, (mountain sorrel), tree hogweed	Up to 2 leaf stage	500 mL	All States		
	Up to 4 leaf stage	750 mL			
Fat hen, field madder, saffron thistle, variegated thistle	Up to 2 leaf stage	500 mL	All States		
	Up to 4 leaf stage	750 mL			
Ox-tongue, wireweed	Up to 2 leaf stage	500 mL	All States		
Fireweed	Up to 2 leaf stage	500 mL	Qld, NSW, ACT, Vic, SA, WA, NT only		
	Up to 4 leaf stage	500 mL			
Common cotula (bird's eye) pheasants eye (adonis)	Up to 4 leaf stage	560 mL	SA only		
	Greater than 4 leaf stage	1.1 L			
Wheat, barley, triticale, cereal Rye	Fumitory	2-6 leaf stage	350 + 200 mL/ha terbutryn (500 g/L)	WA only	
Wheat, barley, triticale, cereal Rye (including undersown with clover and/or lucerne), and these cover crops in vineyards	<b>Suppression of the Following Weeds</b>				
	Dense-flower fumitory	Up to 2 leaf stage	750 mL	All States	
		Up to 4 leaf stage	1.0 L		
	Chickweed, common sowthistle (milk thistle), dock#, hexham scent, (King Island melilot), prickly lettuce, scarlet pimpernel, skeleton weed, sorrel, speedwell, three horned bedstraw, toad rush				
			500 mL-1.0 L		
	Volunteer lupins				
	Crassula (stonecrop)	Up to 5 leaf stage	500 mL		
	Long Storksbill	Up to 4 leaf stage			
	Volunteer field peas	Up to 5 node stage	750 mL		
	Ward's weed	Up to 5 leaf stage	1.0 L		
	Vetch	Up to 2 leaf stage			
	Mouse-eared chickweed			NSW, ACT only	
	Mexican poppy			Qld only	
	Mintweed, spoon cudweed	Up to 4 leaf stage		NSW, ACT only	
	New Zealand spinach	Up to 2 leaf stage	750 mL	Qld only	
	Cleavers	Up to 1 whorl stage	1.0 L	SA only	
	Ball mustard	Up to 4 leaf stage			
	Horehound	Pre-emergence			
Marshmallow	Up to 2 leaf stage				
Wheat, barley, triticale, cereal rye	Wild radish	Up to 4 leaf stage and not more than 120 mm in diameter.	350 mL plus 200 mL MCPA LVE (500 g/L)	WA only	
		Up to 6 leaf stage and not more than 150 mm in diameter.	350 mL plus 200 mL MCPA LVE (500 g/L)	All States	
		Up to 8 leaf stage and not more than 180 mm in diameter.	350 mL plus 200 mL MCPA LVE (500 g/L)		

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHOLDING PERIOD**

**Harvest**

**Cereals, Grapes: NOT REQUIRED WHEN USED AS DIRECTED**

**Grazing:**

**Pasture, Cereals: DO NOT GRAZE OR CUT FOR STOCK FOOD WITHIN 14 DAYS AFTER APPLICATION.**